

Human Journals

Research Article

July 2021 Vol.:19, Issue:1

© All rights are reserved by Jara Gutiérrez et al.

Pilot Project: A Theoretical Framework for The Control of Fertility in A Population Sample of Red Deer from El Monte De El Pardo (Spain)



IJSRM
INTERNATIONAL JOURNAL OF SCIENCE AND RESEARCH METHODOLOGY
An Official Publication of Human Journals



Jara Gutiérrez*^{1,2}, Francisco Javier de Miguel¹

¹Department of Biology of the Autonomous University of Madrid, Darwin 2, 28049 Madrid, Spain.

Submitted: 23 June 2021
Accepted: 29 June 2021
Published: 30 July 2021



www.ijsrm.humanjournals.com

Keywords: Animal Welfare, Contraception, Fertility Control, Immunocontraception, Immunocontraceptive Vaccines, Red Deer

ABSTRACT

This study aims to promote fertility control as a non-lethal means of managing wild animal populations, specifically red deer (*Cervus elaphus*) in a region of Spain. This is the first time that a research plan on the use of immunocontraceptives in red deer in Spain has been laid. This plan is accompanied by a follow-up study to (1) determine the effectiveness of the vaccine over 3 years in the prevention of pregnancy, (2) determine the possible contraindications, (3) identify any weaknesses and drawbacks of the treatment, and (4) propose future improvements. We expect to achieve an average inhibition of reproductive production of more than 80%, and it is hoped that in the first year the effectiveness will reach 90-100%, without notable detrimental effects on the individuals. It is expected that the treated deer will exhibit a secretion response of sufficient anti-GnRH antibodies to achieve an inhibition in their reproductive behaviour and physiology and, therefore, in their fertility. Properly planned use of immunocontraceptive vaccination in red deer populations could prevent overpopulation. Significantly detrimental effects of any kind are not expected in vaccinated animals, and the use of the vaccine does not cause a severe risk to the rest of the fauna and human population. The development of this method is essential in urban and suburban areas, where lethal population control compromises animal welfare, in addition to the potential risk for human populations. We suggest that a greater effort to develop less invasive and non-lethal population control modalities is justifiable.

1. INTRODUCTION

Space constraints can lead to inbreeding and overpopulation for red deer (Patton *et al.* 2007), both having welfare implications. Overpopulation can lead to (1) an increase in the transmission of Lyme disease (Spielman *et al.* 1985, Kilpatrick *et al.* 2014, Stein 2019), (2) a reduction in nutritional condition (Martínez and Hewitt 2001), and (3) an increase in vehicle accidents (Conover *et al.* 1995). Blacklegged ticks (*Ixodes scapularis*) are the main vector of Lyme disease among wild animals and human populations in eastern North America (Barbour & Fish 1993). White-tailed deer (*Odocoileus virginianus*) is a primary host for adult black-legged ticks that feed on the blood of these animals, so the overabundance of deer is a significant determinant of Lyme disease risk (Spielman *et al.* 1985) and a direct relationship between deer numbers and Lyme disease cases has been shown (Kilpatrick *et al.* 2014). A survey revealed that Lyme disease and deer-vehicle collisions were the most prevalent concerns related to deer in suburban environments (Connelly *et al.* 1987, Perry *et al.* 2006).

The general effect of increased inbreeding in animal populations is the increase in the number of homozygous offspring, which means that these individuals can be affected by recessive or deleterious traits (Nabulsi *et al.* 2003). In deer, inbreeding depression is negatively correlated with lifetime and breeding success (Slate *et al.* 2000), therefore, it reduces the ability of populations to survive and reproduce (Jiménez *et al.* 1994). Inbreeding is correlated with brachygnathy (shortened lower jaw) in red deer (Zachos *et al.* 2007).

Recently, debates have arisen with diverse perspectives about the overpopulation management of native ungulates in natural parks (Grignolio *et al.* 2014).

Animal populations that grow beyond certain numbers in ways that conflict with human interests (due to, for instance, disease transmissions and vehicle collisions) are sometimes regulated by humans. This is often done via control over mortality, that is, through hunting, or via control over fertility (Kirkpatrick 1999).

Some national parks have reintroduced predators such as wolves (*Canis lupus*), among other reasons to reduce the number of wild ungulates. However, this practice can be challenged in terms of both direct animal welfare considerations for both the translocated predators and the animals they predate (Horta 2010) and indirect ones as it can generate the opposition of farmers

(Linnell *et al.* 2002) and hunters (Licht *et al.* 2010), who may react by proposing the killing of the reintroduced animals. In addition, they may be harmful to animals without being effective on high-density populations (Chair of national parks *et al.* 2019).

Until very recently, hunting has been the main means of reducing the number of animals of numerous species. However, public pressure to protect deer from hunting has increased in the last two decades as a result of growing animal ethics concerns (Turner *et al.* 1992; Fraser and MacRae 2011). As a result, non-lethal methods have been increasingly supported. In addition, in Monte de El Pardo, red deer populations can be considered as belonging to peri-urban or suburban areas, in which hunting, capture, or poisoning is officially prohibited for human safety reasons. The vicinity of urban and rural centers falls into the category of "security zones," in which the use of hunting weapons is prohibited or is controlled by local regulations (Mesa Gutiérrez 2017).

Recent research (e.g. Bowman, 2011; Gionfriddo *et al.*, 2006, 2011a), has shown the need for new population management tools. An example of this is the translocation of individuals, which is still an expensive and complicated task since it is necessary to ensure an appropriate and effective settlement of the transferred individuals. Mortality rates for relocated deer have been much higher than for resident deer (O'Bryan and McCullough 1985; Craven *et al.* 1998).

For all these reasons, fertility control through contraception in ungulate populations has been the subject of study for almost 40 years (Kirkpatrick and Turner, 1985; Turner *et al.* 1992).

In early contraceptive studies in deer, steroid hormones, primarily diethylstilbestrol (DES, a synthetic estrogen) and melengestrol acetate (AMG, an active progestogen), were administered via injectable implants or orally in food (Bell 1975, Matschke 1977a, 1977b, Roughton 1979). However, implants require capture or immobilization methods that are costly, invasive, and even potentially dangerous to the animal (Turner *et al.* 1992).

There has recently been an increase in research on oral baits for deer in the Mediterranean bush since factors such as season, specificity, and palatability can influence the preferences of individuals. For example, alfalfa bait can be used for oral administration of drugs or vaccines in deer (Martínez-Guijosa *et al.*, 2017). Patton *et al.* (2007) reviewed the possible negative effects of steroids as hormonal contraceptives in ungulates. One of the major disadvantages of these

methods is the fact that they can also act as androgens or antiandrogens, estrogens or antiestrogens, and often have glucocorticoid-like effects, exerting a wide variety of biological effects beyond their contraceptive effects (Patton et al. 2007). Critics object that oral administration of contraceptives can be unreliable, because they can be ingested by individuals who are not the subjects of the study, and they require a frequent, often daily, intake of the steroid (Turner et al. 1992). It has also been argued that it can have other effects that are not desirable because it can affect the behavior of the animals that consume it and can be passed up the food chain (Kirkpatrick and Turner, 1985; Kirkpatrick and Turner 1991).

Immunocontraception is based on the same principle used in the prevention of many diseases: a vaccine made up of proteins that cause the production of antibodies. This means that immunocontraceptive vaccines are based on the animal's ability to produce antibodies against gamete proteins, reproductive hormones, and other proteins essential for reproductive activity (Kirkpatrick et al. 2011). Antibodies that the animal generates interfere with the biological activity of reproductive proteins (Talwar and Gaur 1987) and, depending on the antigen and the vaccine formulation, these can be effective from 1 to 4 years, or even longer, both from single injections and from multiple injections (Miller et al. 1999; Miller et al. 2013).

Numerous immunocontraception procedures interfere with the reproductive process at different levels (e.g. pituitary gland, gonads, sperm, and eggs). The main ones are classified into PZP (Porcine Zona Pellucida) vaccines and GnRH (Gonadotropin-Releasing Hormone) vaccines:

1.1. PZP vaccines (Porcine Zona Pellucida)

PZP is a combination of three zona pellucida (ZP) proteins extracted from pig ovaries. This vaccine causes the female to produce antibodies that interfere with the approach of the sperm to the egg (Florman and Wassarman 1985). After testing PZP-type vaccines in Australian opossums (*Trichosurus vulpecula*) (Duckworth et al., 1999), the Tammar wallaby (*Macropus eugenii*) (Kitchener et al., 2002), koala (*Phascolarctos cinereus*) (Kitchener et al., 2009b), and various eutherian mammals, it was concluded that the native porcine protein formed by glycosylated ZP3 is the most immunogenic of the proteins (Koyama et al., 1996; Kitchener et al., 2002). The effectiveness of the different vaccines tested is 70-80% (Moros 2015). Immunocontraception using PZP-type vaccines generates an inhibition of fertility greater than 95%, with reversible

effects and without affecting females that were already pregnant, and without producing secondary effects on behavior (Kirkpatrick et al. 1990; Kirkpatrick et al. 1991) or noticeable effects on well-being (Killian et al. 2006a). It is not detected in the flesh of the animals (Miller et al. 2013). However, it has some disadvantages. First, inoculation in the female must be performed twice in 4-6 months, with administration accompanied by an adjuvant (a general immunostimulant). Second, unless the population suffers from very high mortality rates, the vaccine itself does not cause immediate or rapid population decline, although it can stabilize a deer population within 2-3 years (Kirkpatrick 1999).

1.2 GnRH vaccines

GnRH, or gonadotropin-releasing hormone, is a hormone that stimulates the release of gonadotropins from the pituitary gland, triggering the cascade of reproductive hormones (estrogen, progesterone, and testosterone) that lead to sperm production in males and ovulation in females (Miller et al. 2013).

Anti-GnRH vaccines stimulate the production of antibodies that bind to the GnRH hormone, that is, antibodies against the individual's GnRH (United States Department of Agriculture 2007). GnRH is a neuropeptide hormone that is naturally secreted from neurons in the hypothalamus and is responsible for stimulating the synthesis and release of two fundamental hormones for proper reproductive physiology: LH (luteinizing hormone) and FSH (follicle-stimulating hormone) in the pituitary gland ("Hazum and Conn, 1988" in Baker et al., 2004). Specifically, long-term treatment with GnRH agonists prevents ovulation, by decreasing GnRH receptors on gonadotropic cells of the adenohypophysis, decreasing the sensitivity of the receptor to GnRH (Nett et al. 1975), decreasing the pituitary LH content, and suppressing the pulsatile secretion of LH and FSH into the bloodstream (Aspden et al. 1996).

To achieve this, these vaccines must be composed of numerous GnRH peptide molecules coupled to a protein (e.g. keyhole limpet hemocyanin [KLH] or blue protein [*Concholepus concholepus*]) thus forming what is known as a "conjugate", and its application in treated animals requires an adjuvant, like PZP (Miller et al. 2013). By binding to GnRH, the antibodies reduce the ability of GnRH to stimulate the release of these sex hormones. As a result,

sexual activity decreases, and animals remain in a non-reproductive state as long as there is a sufficient level of antibody activity (United States Department of Agriculture 2007).

Briefly, immunization against GnRH causes cessation of ovulation and follicular development in females (Patton et al. 2007), through the reduction of the secretion of essential reproductive hormones (Fagerstone et al., 2006; Miller et al., 2008; Miller et al., 2004). The effects on the individual are not lifelong. Following a decrease in antibody titers, natural reversion occurs in most immunized animals (Keeling and Crichton, 1984 in Patton et al., 2007), although the effects can last several years.

Single-dose GnRH vaccines were effective in reducing a population of wild boars for a period of up to 36 weeks when administered to female boars during the reproductive season (Killian et al. 2003), while the response to the vaccine in males was lower (Killian et al., 2006).

This vaccine does not affect non-reproductive behavior or social organization of the species, and the protein antigens are broken down into amino acids in the gastrointestinal tract so that the vaccine does not pass through the food chain (Miller et al. 2013). It has proven to be safe for pregnant females, without interfering with the pregnancy in deer (Miller et al., 2008), elk (*Cervus Canadensis*), American bison (*Bison bison*), and wild horses (*Equus ferus caballus*) (Miller et al. 2004; Killian et al. 2008a; Powers et al. 2012).

Gona ConTM is one of the commercial brands of anti-GnRH vaccines that are most used for the control of fertility in species such as red deer. In fact, in female red deer, a single injection of *GonaConTM* has resulted in infertility for at least 1 year (Hobbs et al. 2000) and up to 2-4 years (United States Department of Agriculture 2007), with 80-100% effectiveness (Killian et al. 2008b). Additionally, *GonaConTM* significantly reduces reproductive behaviors for at least 2 years (Killian et al. 2008b), including heat cycles, which can serve as a tool for controlling the transmission of venereal diseases and diseases transmitted during childbirth (Miller et al. 2013).

In summary, the immunocontraceptive vaccination approach in wild populations is simple to administer, is effective over several years, and has little or no contraindications to treated individuals. The most recent studies show that they are safe and effective in the short term (Killian et al. 2004; Killian et al. 2006a).

For red deer, GnRH-type vaccines seem most suitable. Among all of the animal endocrine suppression modalities developed to date, GnRH immunocontraception probably poses the lowest animal welfare risks (Hampton *et al.* 2015). These vaccines could represent an at least partial solution to the problem of managing overpopulation in deer living near cities (Gionfriddo *et al.* 2006).

2. MATERIALS AND METHODS

2.1 Animals

For this action plan, 200 vaccinated individuals and 50 control individuals will be captured, sedated, identified, and tracked during two consecutive years. All individuals inhabit El Monte de El Pardo (Madrid, Spain, ~16,000 ha). Control animals will be administered a dose equivalent to that of the vaccine, but of a saline solution (Massei *et al.* 2018). The study will begin in 2021 and will have a total duration of 3 years.

As far as possible, the ages and body masses of the treated individuals will be similar, both in the treated group and in the control group (Turner *et al.* 1992). Female individuals of similar reproductive age will be identified following the criteria of scientific guides for hunting species (e.g. Sáenz de Buruaga *et al.*, 2001). Furthermore, the age of each individual will be estimated by observing their dentition, specifically through tooth eruption and wear (Azorit 2011).

2.2 Vaccination

Females of reproductive age (approximately 2-10 years old) will be vaccinated (Carranza 2017). There is no disadvantage in vaccinating pregnant females since the vaccine does not negatively impact the wellbeing of pregnant deer (Miller *et al.*, 2008). Individuals will be vaccinated in July, as has been described in similar studies (Killian *et al.* 2005).

Individuals will be captured and immobilized by tranquilizer darts fired from a distance using a Pneu-Dart rifle (Model 171; Pneu-Dart, Inc., Williamsport, PA), which will contain an anesthesia mixture consisting of Telazol® (4.5 mg/kg; Fort Dodge Animal Health, Fort Dodge, IA, USA) and xylazine hydrochloride (2.5 mg/kg; Bayer, Leverkusen, North Rhine-Westphalia, Germany) (Murray *et al.* 2000; Killian *et al.* 2005; Killian *et al.* 2008b; Gionfriddo *et al.* 2008;

Gionfriddo *et al.* 2011b). Tranquilizer darts will be fired from authorized vehicles traveling on accessible forest roads and trails, or on foot when appropriate.

Once immobilized, individuals will be covered with a thermal blanket and undergo a veterinary examination. The evaluation of the animals will serve to detect possible symptoms or injuries compatible with diseases, wounds, and trauma. The deer will also be equipped with ear tags and radio-tracking collars (Gionfriddo *et al.* 2008).

The body condition of all individuals will be evaluated by palpation of the body contours because it is especially related to fat stores (e.g. Couturier *et al.*, 2009; Finger *et al.*, 1981; Mattiello *et al.*, 2009; Serrano *et al.*, 2008). Body condition will serve as a bioindicator of the health and physical condition of the treated animals.

The formula of *GonaCon*TM administered will contain 1,000 µg of the GnRH peptide combined with AdjuVacTM adjuvant (Miller *et al.*, 2008) as a 1 ml dose injection in the rump (Killian *et al.* 2008b; Gionfriddo *et al.* 2011b). *GonaCon*TM must be injected intramuscularly (United States Department of Agriculture 2007) and it is expected that after a few days the vaccinated animal will begin to generate antibodies.

The inoculation with the vaccine will be administered through a single injection since *GonaCon*TM is capable of inducing contraceptive effects lasting several years with a single dose (Hobbs *et al.* 2000; Killian *et al.* 2008b; Miller *et al.* 2013). The immune response of individuals is variable, with effects lasting between 2 and 5 years (Madrirdejos 2017), or even up to 6 years (Killian *et al.* 2008b).

50 female deer will also be captured but not vaccinated and will serve as an untreated control group (Gionfriddo *et al.* 2008). Control animals will be administered a dose equivalent to that of the vaccine but of a saline solution (Massei *et al.* 2018).

Regular blood samples will be taken from all of the treated individuals, both control and treated groups. Blood samples will be refrigerated and taken to the regional laboratories in Madrid for analysis.

An injection of Tolazine TM (tolazoline: 3.0 - 4.0 mg/kg; Lloyd, Inc., Shenandoah, IA) will be administered to reverse anesthetic effects and to speed the recovery of individuals before release (Gionfriddo *et al.* 2008; Killian *et al.* 2008b; Gionfriddo *et al.* 2011b).

2.3 Tracing

The effectiveness of *GonaCon*TM will be quantified by measuring physiological, ethological and mobility parameters, including the comparison of these data with the results of the control group in a longitudinal study.

The vaccinated individuals will be monitored by discontinuous radio monitoring at discrete or random times during the study period. This technique is useful for determining the range of individuals and their movements (Harris *et al.* 1990).

Periodically every 3-4 months and until the end of the 3-year plan, vaccinated individuals and individuals from the control group will be immobilized again to collect blood samples to quantify progesterone and antibody levels. An evaluation of the general state of their health will be carried out, e.g. observation of possible reactions at the injection site (Massei *et al.* 2018), motor difficulties in the extremities, lameness, abnormal behaviors (Gionfriddo *et al.* 2011b), and body condition (Killian *et al.* 2006a). Additionally, the evaluation of females will include abdominal palpation between April and May to evaluate pregnancies (Miller & Killian 2000).

Behavioral observations will be made 3 times a day at 7:30 a.m., 12:00 p.m., and 4:00 p.m. until mid-January, and then twice a day at 7:30 a.m. and 4:00 pm, at the end of February (Killian *et al.* 2008b). The observation periods will last 30 minutes but will be lengthened in those cases in which reproductive activity is observed, to record as much information as possible for all active individuals. The criteria used to categorize reproductive behaviors will be (1) number of sexual encounters, (2) total time of sexual activity (in days), and (3) number of heat events (Miller & Killian 2000).

2.4 Statistical analysis

Statistical analyzes will be performed comparing the data collected in the vaccinated and control group in the initial vaccination, 12 and 24 months later. Generalized Linear Mixed Model (GLMM) was used to assess the effect of *GonaCon*TM on body condition scores and on factors

that potentially affect the immune response to vaccination with GonaCon™. Treatment group, estimated age, study stage, and pregnancy status at each stage were included as fixed effects. A step-by-step backward selection based on Akaike Information Criteria (AIC) will be used to choose the optimal model (Massei *et al.* 2018).

3. EXPECTED RESULTS

Using a single dose of the *GonaCon™* vaccine, we hope to find the effectiveness of inhibiting the reproduction of females that is greater than 95% in the first year, and greater than 80% in the following two years.

Simultaneously, we expect serum progesterone levels (ng/mL) to be lower in the treated group than in the control group, since the latter will be able to go into heat and have a successful pregnancy. Additionally, the levels of anti-GnRH antibody titers are expected to be higher in the treated group than in the control group, and high enough to block reproductive physiology and consequent reproductive behaviours in vaccinated females (detectable at a dilution of 1:32,000-1:64,000) (Quy *et al.* 2014).

Likewise, a total of almost total absence of unwanted side effects (contraindications) derived from the injection of the vaccine in all or the vast majority of animals is expected. Aspects such as stress behaviours, variations in movement and activity, and reactions in the injection area will be evaluated during the follow-up.

4. DISCUSSION

Our results are expected to indicate that the vaccine is suitable for use in red deer populations to stabilize populations, and can be extrapolated to other species in urban, peri-urban areas and regional parks of the Iberian Peninsula, such as fallow deer and boars. Along with red deer, these populations live relatively close to human populations and are likely to be targeted by humans.

The responsiveness of red deer to fertility control through *GonaCon™* vaccines should encourage researchers and those interested in the control of red deer or other ungulate populations to continue efforts to optimize multi-year contraceptive-inducing vaccines. Recent authors have pointed out the scarcity of scientific studies on the effects of contraceptive vaccines on the physiology and behavior of wild animals (Quy *et al.* 2014).

The effectiveness of the immunocontraceptive vaccine in wild red deer populations will be assessed by medium and long-term follow-ups in both vaccinated individuals and those in the control group.

This population control method avoids the use of lethal force that poses a threat to humans as well the targeted animals. Likewise, controlling the number of red deer present in Monte de El Pardo will reduce the potential threats that derive directly from collisions with vehicles and the transmission of diseases.

We suggest that a greater effort to develop less invasive and non-lethal population management modalities is justifiable. The current theoretical framework allows a first advance for the development of population control that considers the welfare of individual animals.

5. CONCLUSIONS

The implementation of immunocontraception in red deer in Spain will yield results of high interest to the scientific community. In addition, it is expected that stability in the number of deer will reduce potential threats that derive directly from collisions with vehicles and the transmission of diseases.

We have enough scientific evidence to believe that our results will be favorable once they are implemented in practice, which will allow us to support further projects in other territories of Spain, and probably in other countries.

Immunocontraceptive vaccine development improves over the years, and contributing to its advance represents contributing to the welfare of individual animals. For example, the latest scientific advances are moving in the direction of producing an oral species-specific GnRH vaccine bait which would replace the injections, thereby reducing the cost and the risk to the health of the animals (USDA 2007). This approach could reduce the stress of capture and the risk of anesthesia in future *GonaCon*TM vaccination campaigns.

Author's contributions

Jara Gutiérrez (J.G) contributed to the study conception and design, search, compilation, and revision of the literature and writing of the manuscript. Francisco Javier de Miguel (F.J. de M.)

contributed to the supervision of the article and critical revision. All authors read and approved the final manuscript.

Declarations of interest

None.

Funding

This research did not receive any specific funding.

REFERENCES

1. **Aspden WJ, Rao A, Scott PT, Clarke IJ, Trigg TE, Walsh J and D'Occhio MJ** 1996 Direct actions of the luteinizing hormone-releasing hormone agonist, deslorelin, on anterior pituitary contents of luteinizing hormone (LH) and follicle-stimulating hormone (FSH), LH and FSH subunit messenger ribonucleic acid, and plasma concentrations of. *Biology of Reproduction*
2. **Azorit C** 2011 Guía para la determinación de la edad del ciervo ibérico (*Cervus elaphus hispanicus*) a través de su dentición: revisión metodológica y técnicas de elección.. *Anales de la Real Academia de Ciencias Veterinarias de Andalucía Oriental*
3. **Baker DL, Wild MA, Connor MM, Ravivarapu HB, Dunn RL and Nett TM** 2004 Gonadotropin-releasing hormone agonist: A new approach to reversible contraception in female deer. *Journal of Wildlife Diseases*
4. **Barbour AG and Fish D** 1993 The biological and social phenomenon of lyme disease. *Science*
5. **Bell RL and TJP** 1975 Hormone implants control reproduction in white-tailed deer. *Wildlife Society Bulletin*
6. **Bowman JL** 2011 Managing white-tailed deer: Exurban, suburban, and urban environments. *Biology and Management of White-Tailed Deer* p.
7. **Carranza J** 2017 Ciervo – *Cervus elaphus*. En: *Enciclopedia Virtual de los Vertebrados Españoles*. Salvador, A., Barja, I. (Eds.). Museo Nacional de Ciencias Naturales p.
8. **Cátedra de parques nacionales, Red de parques nacionales and Universidad politécnica de Madrid** 2019 Gestión de ungulados silvestres en parques nacionales: 1–55.
9. **Connelly NA, Decker DJ and Wear S** 1987 Public tolerance of deer in a suburban environment: implications for management and control. *Proc. Eastern Wildl. Damage Cont. Conf.* pp. 3:207-218
10. **Conover MR, Pitt WC, Kessler KK, Dubow TJ and Sanborn WA** 1995 Review of human injuries, illnesses, and economic losses caused by wildlife in the United States. *Wildlife Society Bulletin (1973-2006)*
11. **Couturier S, Côté SD, Huot J and Otto RD** 2009 Body-condition dynamics in a northern ungulate gaining fat in winter. *Canadian Journal of Zoology*
12. **Craven S, Barnes T and Kania G** 1998 Toward a professional position on the translocation of problem wildlife. *Wildlife Society Bulletin*
13. **Fagerstone KA, Miller L, A, Bynum K, S, Eisemann JD and Yoder CA** 2006 When, Where and for What Wildlife Species Will Contraception Be a Useful Management Approach?. *Proceedings of the Vertebrate Pest Conference*
14. **Finger SE, Brisbin IL, Smith MH and Urbston DF** 1981 Kidney Fat as a Predictor of Body Condition in White-Tailed Deer. *The Journal of Wildlife Management*
15. **Florman HM and Wassarman PM** 1985 O-linked oligosaccharides of mouse egg ZP3 account for its sperm receptor activity. *Cell*
16. **Gionfriddo J, P, Eisemann JD, Sullivan K, J, Healey R, S and Miller L, A** 2006 Field Test of GonaCon™

Immunocontraceptive Vaccine in Free-Ranging Female White-Tailed Deer. *Proceedings of the Vertebrate Pest Conference*

17. **Gionfriddo J, P, Gates N, B, DeNicola A, J, Fagerstone K, A and Miller L, A** 2008 Field Test of GonaCon™ Immunocontraceptive Vaccine in Free-Ranging Female Fallow Deer. *Proceedings of the Vertebrate Pest Conference*

18. **Gionfriddo JP, Denicola AJ, Miller LA and Fagerstone KA** 2011a Efficacy of GnRH immunocontraception of wild white-tailed deer in New Jersey. *Wildlife Society Bulletin*

19. **Gionfriddo JP, Denicola AJ, Miller LA and Fagerstone KA** 2011b Health effects of GnRH immunocontraception of wild white-tailed deer in New Jersey. *Wildlife Society Bulletin*

20. **Grignolio S, Heurich M, Šprem N and Apollonio M** 2014 The management of ungulates in protected areas. *Behaviour and Management of European Ungulates*: 178–191.

21. **Hampton JO, Hyndman TH, Barnes A and Collins T** 2015 Is wildlife fertility control always humane?

22. **Harris S, Cresswell WJ, Forde PG, Trewhella W, Woollard TH and Wray S** 1990 Home range analysis using radio-tracking data? A review of problems and techniques particularly as applied to the study of mammals. *Mammal Review* 20: 97–123.

23. **Hazum E and Conn PM** 1988 Molecular mechanism of gonadotropin releasing hormone (gnrh) action. I. The GNRH receptor. *Endocrine Reviews*

24. **Hobbs NT, Bowden DC and Baker DL** 2000 Effects of Fertility Control on Populations of Ungulates: General, Stage-Structured Models. *The Journal of Wildlife Management*

25. **Horta O** 2010 The Ethics of the Ecology of Fear against the Nonspeciesist Paradigm: A Shift in the Aims of Intervention in Nature. *Between the Species: An Online Journal for the Study of Philosophy and Animals*

26. **Jiménez JA, Hughes KA, Alaks G, Graham L and Lacy RC** 1994 An experimental study of inbreeding depression in a natural habitat. *Science*

27. **KEELING BJ and CRIGHTON DB** 1984 REVERSIBILITY OF THE EFFECTS OF ACTIVE IMMUNIZATION AGAINST LH-RH. *Immunological Aspects of Reproduction in Mammals* p.

28. **Killian G, Eisemann J, Wagner D, Werner J, Shaw D, Engeman R and Miller L, A** 2006a Safety and Toxicity Evaluation of GonaCon™ Immunocontraceptive Vaccine in White-Tailed Deer. *Proceedings of the Vertebrate Pest Conference*

29. **Killian G, Miller L, Rhyan J, Dees T, Perry D and Doten H** 2003 Evaluation of GnRH contraceptive vaccine in captive feral swine in Florida. *Proceedings of the 10th Wildlife Damage Management Conference*

30. **Killian G, Miller L, Rhyan J and Doten H** 2006b Immunocontraception of florida feral swine with a single-dose GnRH vaccine. *American Journal of Reproductive Immunology*

31. **Killian G, Miller LA, Diehl NK, Rhyan J, Thain DS, Miller LA, Rhyan J, Thain DS, Diehl NK, Rhyan J, Thain DS, Miller LA, Rliyan J and Thaiii D** 2004 Evaluation of three contraceptive approaches for population control of wild horses. *Proceedings of the 21st Vertebrate Pest Conference*

32. **Killian G, Thain D, Diehl NK, Rhyan J and Miller L** 2008a Four-year contraception rates of mares treated with single-injection porcine zona pellucida and GnRH vaccines and intrauterine devices. *Wildlife Research* p.

33. **Killian G, Wagner D, Fagerstone K and Miller L** 2008b Long-Term Efficacy and Reproductive Behavior Associated with GonaCon Use in White-Tailed Deer (*Odocoileus virginianus*). *Proceedings of the Vertebrate Pest Conference*

34. **Killian G, Wagner D and Miller L** 2005 Observations on the Use of the GnRH Vaccine Gonacon™ in Male White-Tailed Deer (*Odocoileus virginianus*). *Proceedings of the 11th Wildlife Damage Management Conference*

35. **Kilpatrick HJ, Labonte AM and Stafford KC** 2014 The relationship between deer density, tick abundance, and human cases of lyme disease in a residential community. *Journal of Medical Entomology*

36. **Kirkpatrick JF** 1999 Control de la fertilidad en la vida salvaje: un nuevo paradigma para el tratamiento humano de los animales Author (s): Jay F . Kirkpatrick Source : Teorema : Revista Internacional de Filosofia , 1999 , Vol . 18 , No . 3 , Los derechos de Published by : 18: 137–148.

37. **Kirkpatrick JF, Liu IK and Turner JW** 1990 Remotely delivered immunocontraception in feral horses. *Wildlife Society Bulletin*

38. **Kirkpatrick JF, Liu IM, Turner JW and Bernoco M** 1991 Antigen recognition in feral mares previously immunized with porcine zona pellucidae.. *Journal of reproduction and fertility. Supplement*
39. **Kirkpatrick JF, Lyda RO and Frank KM** 2011 Contraceptive Vaccines for Wildlife: A Review
40. **Kirkpatrick JF and Turner, JW** 1985 Chemical Fertility Control and Wildlife Management. *BioScience*
41. **Kirkpatrick JF and Turner JW** 1991 Reversible Contraception in Nondomestic Animals. *Source: Journal of Zoo and Wildlife Medicine Journal of Zoo and Wildlife Medicine*
42. **Licht DS, Millspaugh JJ, Kunkel KE, Kochanny CO and Peterson RO** 2010 Using small populations of wolves for ecosystem restoration and stewardship. *BioScience*
43. **Linnell JDC, Andersen R, Andersone Ž, Balčiauskas L, Blanco J, Boitani L, Brainerd S, Breitenmoser U, Kojola I, Liberg O and Others** 2002 The fear of wolves: a review of wolf attacks on humans. *Norw. Inst. Nature Res. Oppdragsmelding*
44. **Madrídejos A** 2017 Una vacuna contraceptiva para frenar la superpoblación de jabalís. <https://www.elperiodico.com/es/medio-ambiente/20170427/jabali-matadepera-inicio-vacuna-contraceptiva-uab-6001927>
45. **Martínez-guijosa J, Casades-martí L, Royo-hernández L, Gortázar C, Aranaz A, González-Barrio D and Ruiz-Fons F** 2017 Ensayo de campo de cebos orales para ciervo (*Cervus elaphus*) en monte mediterráneo. XIII Congreso SECEM, 6-9 diciembre, Guadalajara, España
46. **Martínez M. A and Hewitt D** 2001 Sobre población de venado cola blanca en el noreste de México. *Ciencia UANL* 4: 177–181.
47. **Massei G, Koon KK, Law SI, Gomm M, Mora DSO, Callaby R, Palphramand K and Eckery DC** 2018 Fertility control for managing free-roaming feral cattle in Hong Kong. *Vaccine*
48. **Matschke GH** 1977a Microencapsulated Diethylstilbestrol as an Oral Contraceptive in White-Tailed Deer. *The Journal of Wildlife Management*
49. **Matschke GH** 1977b Fertility Control in White-Tailed Deer by Steroid Implants. *The Journal of Wildlife Management*
50. **Mattiello S, Andreoli E, Stefanelli A, Cantafora A and Bianchi A** 2009 How to evaluate body conditions of red deer (*Cervus elaphus*) in an alpine environment? . *Italian Journal of Animal Science*
51. **Mesa Gutiérrez MJ** 2017 Marco penal y administrativo de la caza y responsabilidad civil en derecho español. Complutense University of Madrid
52. **Miller L, Gionfriddo J, Rhyan J, Fagerstone K, Wagner D and Killian G** 2008a GnRH immunocontraception of male and female white-tailed deer fawns. *Human-Wildlife Conflicts*
53. **Miller L and Killian G** 2000 Seven years of white-tailed deer immunocontraceptive research at Penn State University: A comparison of two vaccines. *Proceedings of the 9th Wildlife Damage Management Conference*
54. **Miller LA, Fagerstone KA and Eckery DC** 2013 TWENTY YEARS OF IMMUNOCONTRACEPTIVE RESEARCH: LESSONS LEARNED. *Journal of Zoo and Wildlife Medicine*
55. **Miller LA, Gionfriddo JP, Fagerstone KA, Rhyan JC and Killian GJ** 2008b The single-shot GnRH immunocontraceptive vaccine (GonaCon™) in white-tailed deer: Comparison of several GnRH preparations. *American Journal of Reproductive Immunology*
56. **Miller LA, Johns BE and Killian GJ** 1999 Long-term effects of PZP immunization on reproduction in white-tailed deer. *Vaccine*
57. **Miller LA, Rhyan JC and Drew M** 2004 Contraception of bison by GnRH vaccine: A possible means of decreasing transmission of brucellosis in bison. *Journal of Wildlife Diseases*
58. **Moros C** 2015 Análisis Molecular, Proteómico y Filogenético de la Zona Pelúcida de Mamíferos. Universidad de Murcia
59. **Murray S, Monfort SL, Ware L, McShea WJ and Bush M** 2000 Anesthesia in female white-tailed deer using Telazol® and xylazine. *Journal of Wildlife Diseases*
60. **Nabulsi MM, Tamim H, Sabbagh M, Obeid MY, Yunis KA and Bitar FF** 2003 Parental consanguinity and congenital heart malformations in a developing country. *American Journal of Medical Genetics*
61. **Nett TM, Akbar AM and Phemister RD** 1975 Levels of luteinizing hormone, estradiol and progesterone in

serum during the estrous cycle and pregnancy in the Beagle bitch. *Proceedings of the Society for Experimental Biology and Medicine*

62. **O'Bryan MK and McCullough DR** 1985 Survival of Black-Tailed Deer Following Relocation in California. *The Journal of Wildlife Management*

63. **Patton ML, Jöchle W and Penfold LM** 2007 Review of contraception in ungulate species

64. **Perry K, R, Arjo W, M, Bynum K, S and Miller L, A** 2006 GnRH Single-Injection Immunocontraception of Black-Tailed Deer. *Proceedings of the Vertebrate Pest Conference*

65. **Powers JG, Baker DL, Ackerman MG, Bruemmer JE, Spraker TR, Conner MM and Nett TM** 2012 Passive transfer of maternal GnRH antibodies does not affect reproductive development in elk (*Cervus elaphus nelsoni*) calves. *Theriogenology*

66. **Quy RJ, Massei G, Lambert MS, Coats J, Miller LA and Cowan DP** 2014 Effects of a GnRH vaccine on the movement and activity of free-living wild boar (*Sus scrofa*). *Wildlife Research*

67. **Roughton RD** 1979 Effects of Oral Melengestrol Acetate on Reproduction in Captive White-Tailed Deer. *The Journal of Wildlife Management*

68. **Sáenz de Buruaga M, Lucio AJ and PURROY J** 2001 Reconocimiento de sexo y edad en especies cinegéticas. Edileisa, León

69. **Serrano E, Alpizar-Jara R, Morellet N and Hewison AJM** 2008 A half a century of measuring ungulate body condition using indices: Is it time for a change?. *European Journal of Wildlife Research*

70. **Slate J, Kruuk LEB, Marshall TC, Pemberton JM and Clutton-Brock TH** 2000 Inbreeding depression influences lifetime breeding success in a wild population of red deer (*Cervus elaphus*). *Proceedings of the Royal Society B: Biological Sciences*

71. **Spielman A, Wilson ML, Levine JF and Piesman J** 1985 Ecology of *Ixodes dammini*-borne human babesiosis and Lyme disease.. *Annual review of entomology. Vol. 30*

72. **Stein RA** 2019 Lyme disease. *Encyclopedia of Environmental Health* p.

73. **Talwar GP and Gaur A** 1987 Recent developments in immunocontraception. *American Journal of Obstetrics and Gynecology*

74. **Turner JW, Liu IKM and Kirkpatrick JF** 1992 Remotely Delivered Immunocontraception in Captive White-Tailed Deer. *The Journal of Wildlife Management*

75. **United States Department of Agriculture A and PHIS** 2007 GonaCon™—Birth Control for Deer: Questions and Answers

76. **USDA** 2007 GonaCon™ – Birth Control for Deer : Questions and Answers. *Wildlife Research*

77. **Zachos FE, Althoff C, Steynitz Y V., Eckert I and Hartl GB** 2007 Genetic analysis of an isolated red deer (*Cervus elaphus*) population showing signs of inbreeding depression. *European Journal of Wildlife Research*